

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Naturalure[®]

Fruit Fly Bait Concentrate

INSECTICIDE

ACTIVE CONSTITUENT: 0.24 g/L SPINOSAD

GROUP	5	INSECTICIDE
-------	---	-------------

For the control of Fruit flies, including Queensland fruitfly and Medfly, infesting various tree, fruit, nut, vine and vegetable crops and ornamentals and on fly resting sites, including non-crop vegetation.

Pack size: 4 L, 5 L, 10 L, 20 L, 100 L, 208 L, 1000 L

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre.
Phone: *Australia* 13 11 26. *New Zealand* 0800 764 766.

**EMERGENCY RESPONSE
(ALL HOURS)**
RING FROM ANYWHERE IN
AUSTRALIA
1800 370 754
(LOCAL CALL FEE ONLY)

SAFETY DIRECTIONS

- May irritate the eyes.
- Avoid contact with eyes.
- If product is in eyes, wash it out immediately with water.
- After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

IN A TRANSPORT
EMERGENCY ONLY
DIAL 000
FOR POLICE OR
FIRE BRIGADE



Naturalure[®] Fruit Fly Bait Concentrate contains proteins from a plant extract to which some people may be allergic. If you are concerned about a potential allergic response, it is advisable to wear gloves and/or some face protection such as a dust mask when mixing or using the product. Take care to avoid contact of the product or spray solution with skin and eyes.

SAFETY DATA SHEET

Additional information is listed on the Safety Data Sheet for **NATURALURE[®] FRUIT FLY BAIT CONCENTRATE** which is available from Corteva Agriscience on request. Call Customer Service Toll Free on 1-800 700 096 or visit www.corteva.com.au



[™] © Trademarks of Corteva Agriscience and its affiliated companies.

Visit us at Corteva.com.au

DIRECTIONS FOR USE

SITUATION	PEST	USE RATE	CRITICAL COMMENTS
Tree, fruit, nut, vine and vegetable crops and ornamentals	Fruit flies, including Queensland fruit fly (<i>Bactrocera tryoni</i>) & Mediterranean fruit fly (<i>Ceratitis capitata</i>)	1 L/ha	Mix 1 part of Naturalure® Fruit Fly Bait Concentrate with 6.5 parts of water. Spray solution can be applied as a band or spot spray (see Application section)
Non-crop vegetation and other fruit fly resting sites (for use in area-wide fruit fly eradication or control programs)			Dilute as above or use as a concentrate spray by mixing 1 part of Naturalure® Fruit Fly Bait Concentrate with 1.5 parts of water and apply as a spot spray (see Application section).

AVOID spraying the fruit as phytotoxicity may occur. Mangoes and pears are particularly susceptible to blemishes, discolouration or severe spotting (superficial on the skin only) from protein-based baits. The risk of phytotoxicity is increased during hot, dry conditions and re-application of spray to the same location on the plants. If using a weekly spray program, alternate tree or crop rows to which the product is applied. Re-apply if rain washes bait off the foliage.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION

WITHHOLDING PERIOD: All crops

NOT REQUIRED when used as directed

GENERAL INSTRUCTIONS

Naturalure® Fruit Fly Bait Concentrate is a protein and sugar-based bait containing the active ingredient spinosad. Once applied, fruit flies can detect the bait from several metres away. Naturalure® Fruit Fly Bait Concentrate can be diluted with water (see **MIXING** section below) to produce either a concentrated or a dilute solution. Use the concentrated solution if application equipment will allow, as this will maximise bait longevity and improve rainfastness. If necessary, remove filters and strainers from spraying equipment to avoid blockages.

Note: Baiting for fruit fly control should be supplemented by good orchard management practices such as removal of fallen and/or rotting fruit.

INSECTICIDE RESISTANCE WARNING

GROUP	5	INSECTICIDE
-------	---	-------------

For insecticide resistance management Naturalure® Fruit Fly Bait Concentrate is a Group 5 insecticide. Some naturally occurring insect biotypes resistant to Naturalure® Fruit Fly Bait Concentrate and other Group 5 insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Naturalure® Fruit Fly Bait Concentrate or other Group 5 insecticides are used repeatedly. The effectiveness of Naturalure® Fruit Fly Bait Concentrate on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Corteva Agriscience Australia Limited accepts no liability for any losses that may result from the failure of Naturalure® Fruit Fly Bait Concentrate to control resistant insects

MIXING

Once mixed, constant agitation of the spray solution is recommended to ensure uniformity of spray mixture. Allow agitation system to operate for at least 5 minutes before applying. Once diluted, the solution should be used within 24 hours.

Concentrated Solution: Mix 1 part of Naturalure® Fruit Fly Bait Concentrate with 1.5 parts of water. First add water (equivalent to one-half of the volume of Naturalure® Concentrate to be mixed) to the spray tank and start the agitation system. Then add the full amount of Naturalure® Concentrate followed by an equal amount of water.

Dilute Solution: Mix 1 part of Naturalure® Fruit Fly Bait Concentrate with 6.5 parts of water. First add water (equivalent to the volume of Naturalure® Concentrate to be mixed) to the spray tank or premixing tank and start the agitation system. Then add the full amount of Naturalure® Concentrate followed by the remaining amount of water.

APPLICATION

Begin applications as soon as monitoring traps indicate flies are present and fruit is at a susceptible stage. Repeat applications every 7 days, re-applying sooner if rain washes off the deposit. A large spray droplet size of 4000 to 6000 μ (4-6 mm) is recommended to optimise duration of bait's attractiveness to flies. Apply as either a band or spot spray to the lower canopy of fruiting plants as follows:

Band sprays (concentrated or dilute solution): Apply as a coarse spray in a 1m wide band to the skirt of trees. Apply to one side of every row or every second row of trees.

Spot sprays of the concentrated solution: Apply to trees and foliage as coarse spots of 20 mL per spot of 1m². Apply 125 spots per hectare.

Spot sprays of the dilute solution: Apply to trees and foliage as coarse spots of 50 mL per spot of 1m². Apply 150 spots per hectare.

Spots should be distributed evenly throughout an orchard to optimise effectiveness. Adjust the application of the spots to suit the number of trees per hectare, but do not exceed the application rate above.

CLEANING SPRAY EQUIPMENT

After using Naturalure® Fruit Fly Bait Concentrate empty the tank and completely drain the system. Rinse the tank, pumps, lines, hoses, filters and nozzles by circulating clean water through the system. Drain and repeat the rinsing procedure twice.

PROTECTION OF LIVESTOCK

Dangerous to bees. Avoid direct application or drift of the spray mix onto beehives.

DO NOT spray on plants in flower if bees are active. Once the spray deposit has dried, foraging bees will not be affected.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT allow the product or used containers to enter dams, ponds, waterways or drains.

DO NOT apply in strong winds or any other conditions that may result in drift onto adjacent pastures, crops or water supplies.

PROTECTION OF NON-TARGET INSECTS

Beneficial insects contribute to control of pest outbreaks. Applications of Naturalure® Fruit Fly Bait Concentrate are unlikely to affect lacewings (*Chrysopa* spp.), predatory bugs (*Geocoris*, *Orius* and *Nabis* spp.), spiders and most species of ladybird beetles (*Coccinella*, *Diomus* and *Harmonia* spp.). However some species of beneficial insects may be sensitive to Naturalure® Fruit Fly Bait Concentrate and its use may temporarily reduce populations of parasitoid wasps (especially *Trichogramma* spp.), ants, some beetles and tachinid flies. This may lead to some disruption of IPM systems based on these species, but generally populations will recover.

STORAGE AND DISPOSAL

Keep out of reach of children. Store in the closed, original container in a dry, cool, well-ventilated area out of direct sunlight.

4 L, 5 L, 10 L, 20 L containers

This container can be recycled if it is clean, dry, free of visible residues and has the *drumMUSTER* logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate by adding to the spray tank. Do not dispose of undiluted chemicals on site. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *drumMUSTER* collection or similar container management site. The cap should not be replaced but may be taken separately. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. DO NOT burn empty containers or product.

100 L, 208 L, 1000 L containers

Do NOT remove or tamper with the dry valves or security seal. DO NOT contaminate the drum with water or any other foreign matter. Empty contents fully into application equipment. After each use of the product ensure that the dry valve coupler, delivery system and hoses are disconnected, triple rinsed with clean water and drained. Add the rinsings to the spray tank. When the drum is empty remove the dry valve coupler and return to the point of purchase. The drum remains the property of Corteva Agriscience and must be returned.

SMALL SPILL MANAGEMENT

Sweep up material and contain in a refuse vessel for disposal in the same manner as for containers (see STORAGE and DISPOSAL section).

APVMA Approval Number: 58234/57203

Corteva Agriscience Australia Limited A.B.N. 24 003 771 659

67 Albert Avenue, Chatswood, NSW 2067

www.corteva.com.au

CUSTOMER SERVICE TOLL FREE

1-800 700 096



GMID

DOM/Batch No.:

This product is GHS compliant. No additional GHS hazard and precautionary statements are required under the WorkSafe Australia exemptions for AgVet products.